



What is Agrizest?

Agrizest is a unique Kiwi technology proven to increase vine health and orchard gate returns (OGR) in New Zealand orchards for over 10 years.

Agrizest elicits the production of phenylpropanoids - which are bioactive molecules that are responsible for a range of plant characteristics. These result in:

- Improved vine health
- Reduced damage and stress caused by pest, disease & environmental factors
- Improved fruit quality & yield
- Increased OGR

What is Primazest?

Primazest is specifically formulated to reduce stress from copper application and to support energy production during early growth stages before leaves are fully developed.

Using **Primazest** encourages:

- Healthy growth
- Reduced damage
- Recovery from stress caused by variable spring conditions such as high winds and frost





ProvenResults

Agrizest and Primazest are commercially proven, extensively peer reviewed (data available), independently verified by the scientific community - with data and case studies from the last 10 years.

Agrizest and **Primazest** are authorised by MPI (under the ACVM Act 1997) as exempt from registration and can be applied to all crops with nil withholding and nil waiting period.

Agrizest and Primazest are listed on the Zespri Allowed Other Compounds (AOC) list.



ProvenPerformance

- Increase vine productivity
- Improve kiwifruit quality



ProvenReturns

Just 4 applications of **Agrizest** result in a minimum increase in orchard gate returns of \$5000 per hectare. For example:

	Control	Agrizest	
Bins/Hectare	139	151	9.00%
Class 1 trays/ hectare (Kg)	11054	11954	8.00%
Orchard Gate Returns	\$37,844	\$44,470	\$6,626

Healthier Vines

Better Fruit

Sweeter Returns

Agrizest and Primazest Spray Programme: Kiwifruit

4 Agrizest sprays must be applied on time to maximise improvements in both yield and quality



PRIMAZEST

starting at the 2-leaf stage (qolfball size).



AGRIZEST

prays 1&2 begin 2-3 weeks before flowering.



prays 3&4 begin immediately after the bees are removed. AGRIZEST



after flowering

2-leaf stage

2-3 weeks before flowering

flowering



1st PRIMAZEST SPRAY

applied at the 2 leaf stage

(around 4cm).

applied 2-3 weeks before flowering.




2nd PRIMAZEST SPRAY

time before the 1st Agrizest spray

applied 7 days later (if there is

applied 3-7 days later.



3rd AGRIZEST SPRAY

applied immediately after the bees are removed. 4th AGRIZEST SPRAY

applied 3-7 days later.



Agrizest and Primazest are included in the Zespri Allowed Other Compounds List Agrizest and Primazest are BioGro certified inputs for organic production.



Call us now and get the best out of your orchard this year.

Call 0274633442 or email info@zestbiotech.co.nz





Agrizest and Primazest are safe to use and are authorised by MPI (under the ACVM Act 1997) as exempt from registration. Agrizest and Primazest can be applied on all crops with nil withholding period and nil waiting period.

- No wetting agent required.
- Water rates: dependent on sprayer and canopy -ensure product application rate is 1L per hectare with typical water rates of 500-1500L per hectare.
- Agrizest and Primazest are compatible with most commonly used orchard sprays.
- Compatibility with additional products must be tested prior to application.
- Do not mix with copper or hormones including Actigard and CPPU.





Call us now to get the best out of your orchard this year.

Call 0274633442 or 092383893